

Local algebra bundles and α -jets of mappings.

MARGHERITA BARILE*
FIORELLA BARONE*
WŁODZIMIERZ M. TULCZYJEW**

*Dipartimento di Matematica, Università di Bari, Italy.

**Dipartimento di Fisica, Università di Camerino, Italy.

Abstract. We present the functor associated with a local algebra bundle and the differential structure of the double fibre bundle it produces when applied to a differential manifold.

KEYWORDS: *Local algebras, double bundles, jet bundle functors.*

A.M.S. CLASSIFICATION: 13H10, 58A20

1. Introduction and preliminaries.

Weil algebras and related functors [14] have been quite extensively studied by a number of authors over the past decades ([2], [3], [4], [5], [7], [8], [12]). In the present note we generalize the method of associating a functor with a single local algebra. We start with a local algebra bundle $\alpha : A \rightarrow M$ on a differential manifold M and define a functor which associates with a differential manifold P a double fibre bundle

$$\begin{array}{ccc} & AP & \\ \alpha_P^s \swarrow & & \searrow \alpha_P^t \\ M & & P \end{array} \quad (1)$$

The elements of AP are called α -jets on P .

In the first part of the paper, using some of the main results proved in [1] and [2], we give the proof of the differential structure of (1).

In the second part we present a straightforward proof of the connection between the jets of mappings of Ehresmann type ([6], [7], [9], [10], [11], [13]) and a special class of α -jets.


A list of few basic definitions and conventions now follows.

Differential manifolds will be assumed to be modelled on affine spaces and we will refer to the differential calculus on affine spaces presented in [1].

Let M , P , Q and R be differential manifolds.

Definition 1. A double bundle with total space Q , bases M and P and standard fibre R is a

diagram




such that

- (i) $\sigma : Q \rightarrow M$ and $\tau : Q \rightarrow P$ are surjective submersions;
- (ii) for each $x \in M$ and $p \in P$, there are a chart $(\mathcal{U}, \xi)$ around x , a chart $(\mathcal{W}, \eta)$ around p and a diffeomorphism


$$\chi: \sigma^{-1}(\mathcal{U}) \cap \tau^{-1}(\mathcal{W}) \longrightarrow \xi(\mathcal{U}) \times \eta(\mathcal{W}) \times R$$

such that the following double local trivialization diagram



is commutative.

A double bundle of the form



is called a *trivial double bundle*.

Definition 2. A *local algebra bundle* (l.a.b.) is a (locally trivial) fibre bundle

$$\begin{array}{c} A \\ \downarrow \alpha \\ M \end{array}$$

with a local algebra as its standard fibre.

For each $x \in M$, $A_x = \alpha^{-1}(x)$ is called the *local algebra over x* . If I_{A_x} denotes the only maximal ideal of A_x , we identify $\mathbb{R}$ with $\mathbb{R}1_{A_x}$, so that $A_x = \mathbb{R} + I_{A_x}$. Furthermore we denote by 0_{A_x} the only epimorphism from A_x onto $\mathbb{R}$.

A *local algebra bundle morphism* is a bundle morphism which induces a local algebra morphism on each fibre.

2. The functor associated with a local algebra bundle.

Let

$$\begin{array}{c} A \\ \downarrow \alpha \\ M \end{array} \quad (2)$$

be a local algebra bundle.

Let P be a differential manifold, and consider the set $A_{(x,\cdot)}P$ of all $\mathbb{R}$ -algebra morphisms from $C^\infty(P)$ to A_x which preserve the identity. We set

$$AP = \bigcup_{x \in M} A_{(x,\cdot)}P.$$

The elements of AP are called α -jets on P .

We denote by

$$\alpha_P^s : AP \longrightarrow M$$

the natural surjective mapping (*source*) such that for all $x \in M$,

$$(\alpha_P^s)^{-1}(x) = A_{(x,\cdot)}P.$$

Let $u \in AP$ and $x = \alpha_P^s(u)$. There is just one point $p \in P$ such that the composition $0_{A_x} \circ u$ is the evaluation mapping ev_p at p (cf., e.g., [7], p.296). So we have another surjective mapping (*target*)

$$\alpha_P^t : AP \longrightarrow P$$

such that for all $p \in P$,

$$(\alpha_P^t)^{-1}(p) = A_{(\cdot,p)}P = \bigcup_{x \in M} \{u \in A_{(x,\cdot)}P : 0_{A_x} \circ u = ev_p\}. \quad (3)$$

We set

$$A_{(x,p)}P = A_{(x,\cdot)}P \cap A_{(\cdot,p)}P.$$

Then

$$A_{(x,\cdot)}P = \bigcup_{p \in P} A_{(x,p)}P \quad , \quad A_{(\cdot,p)}P = \bigcup_{x \in M} A_{(x,p)}P$$

It is easy to prove the following

Proposition 3.

(i) To a l.a.b. $\alpha : A \rightarrow M$ corresponds a covariant functor

$$\begin{array}{ccc} & A & \\ \alpha^s \swarrow & & \searrow \alpha^t \\ M & & I \end{array} \quad (4)$$

which associates with any differential manifold P the diagram

$$\begin{array}{ccc}
 & AP & \\
 \alpha_P^s \swarrow & & \searrow \alpha_P^t \\
 & M & P
 \end{array} \tag{5}$$

and with any differentiable mapping $\varphi : P \rightarrow Q$ the diagram

$$\begin{array}{ccccc}
 AP & \xrightarrow{A\varphi} & AQ & & \\
 \alpha_P^s \swarrow & & \searrow \alpha_Q^t & & \\
 & M & P & \xrightarrow{\varphi} & Q \\
 \alpha_P^s \swarrow & & \searrow \alpha_Q^s & & \\
 & M & & & M
 \end{array}$$

where, for all $u \in AP$,

$$A\varphi(u) = u \circ \varphi^*, \tag{6}$$

i.e., for all $f \in C^\infty(Q)$,

$$A\varphi(u)(f) = u(f \circ \varphi).$$

(ii) To a l.a.b. morphism

$$\begin{array}{ccc}
 A & \xrightarrow{\kappa} & B \\
 \alpha \downarrow & & \downarrow \beta \\
 M & \xrightarrow{\mu} & N
 \end{array}$$

corresponds the functor

$$\begin{array}{ccccc}
 & A & & & \\
 \alpha^s \swarrow & & \searrow \alpha^t & & \\
 & M & & I & \\
 \mu \swarrow & & \searrow \kappa & & \\
 & N & & B & \\
 & & \beta^s \swarrow & & \searrow \beta^t \\
 & & & & I
 \end{array}$$


which associates with a differential manifold P the diagram

$$\begin{array}{ccccc}
 & AP & & & \\
 \alpha_P^s \swarrow & & \searrow \alpha_P^t & & \\
 & M & & P & \\
 \mu \swarrow & & \searrow \kappa_P & & \\
 & N & & BP & \\
 & & \beta_P^s \swarrow & & \searrow \beta_P^t \\
 & & & & P
 \end{array}$$

where

$$\kappa_P(u) = \kappa \circ u,$$

and with a differentiable mapping $\varphi : P \rightarrow Q$ the diagram



In this section we will prove that diagram (5) is a double bundle.

To this end, we first consider the functor associated with a trivial l.a.b. over an affine space and apply it to another affine space. The resulting trivial double bundle will be the model for the structure we are looking for.

In the sequel, E will denote an affine space modelled on a vector space X and G an affine space modelled on a vector space Z .

Lemma 4. *Let*

$$\begin{array}{c} E \times \bar{A} \\ \downarrow \varepsilon = pr_1 \\ E \end{array} \quad (7)$$

be a trivial l.a.b. over the affine space E . The corresponding functor, when applied to the affine space G , yields a trivial double bundle with an affine space as its total space.

Proof. By definition, when we apply the functor associated with l.a.b. (7) to G , we get

$$\begin{array}{c} (E \times \bar{A})G \\ \swarrow \varepsilon_G^s \quad \searrow \varepsilon_G^t \\ E \qquad \qquad G \end{array} \quad (8)$$

Then we have

$$\begin{aligned} (E \times \bar{A})G &= \bigcup_{x \in E} (\varepsilon_G^s)^{-1}\{x\} \\ &= \bigcup_{x \in E} (\{x\} \times \bar{A})G \\ &= E \times \bar{A}G. \end{aligned} \quad (9)$$

We have proved the following identification (cf. [2], Prop.9)

$$\bar{A}G = G \times (I_{\bar{A}} \otimes Z), \quad (10)$$

where $I_{\bar{A}}$ is the maximal ideal of $\bar{A}$. As a consequence we have

$$(E \times \bar{A})G = E \times G \times (I_{\bar{A}} \otimes Z), \quad (11)$$

so that $(E \times \bar{A})G$ is an affine space and diagram (8) becomes the following trivial double bundle

$$\begin{array}{ccc} & E \times G \times (I_{\bar{A}} \otimes Z) & \\ pr_1 \swarrow & & \searrow pr_2 \\ & E & G \end{array}$$

■

Let us notice that equalities (10) and (11) still hold if E and G are replaced by open subsets owing to the local character of α -jets which we briefly recall.

Let $\mathcal{W}$ be an open subset of P . From the action of the functor (4) on the inclusion mapping

$$\iota_{\mathcal{W}} : \mathcal{W} \rightarrow P$$

we obtain the mapping

$$A_{\iota_{\mathcal{W}}} : A\mathcal{W} \longrightarrow AP$$

whose restriction to any fibre $A_{(\cdot, p)}\mathcal{W}$ of α_P^t

$$A_p \iota_{\mathcal{W}} : A_{(\cdot, p)}\mathcal{W} \longrightarrow A_{(\cdot, p)}P$$

is a bijection (cf. [2], Sec.4.2). As a consequence, owing to the identification

$$A_{(\cdot, p)}\mathcal{W} = A_{(\cdot, p)}P,$$

we have that $A\mathcal{W}$ is a subset of AP (local character of α -jets) and that $A_{\iota_{\mathcal{W}}}$ is the inclusion mapping $\iota_{A\mathcal{W}}$. This is illustrated by the following diagram

$$\begin{array}{ccccc} & A\mathcal{W} & \xrightarrow{\iota_{A\mathcal{W}}} & AP & \\ \alpha_{\mathcal{W}}^s \swarrow & & \searrow \alpha_{\mathcal{W}}^t & & \searrow \alpha_P^t \\ & \mathcal{W} & \xrightarrow{\iota_{\mathcal{W}}} & P & \\ \downarrow & & \downarrow & & \\ M & \xlongequal{\quad} & M & & \end{array}$$

The following step is to apply the functor associated with the l.a.b. morphism given by a trivialization of (2) to a chart of differential manifold P . The result will be a chart of AP .

In the sequel, M will denote a differential manifold modelled on the affine space E and P a differential manifold modelled on the affine space G .

Let

$$\begin{array}{ccc}
 A_\xi = \alpha^{-1}(\mathcal{U}) & \xrightarrow{\xi^A} & \xi(\mathcal{U}) \times \bar{A} \\
 \alpha \downarrow & & \downarrow \varepsilon = pr_1 \\
 \mathcal{U} & \xrightarrow{\xi} & \xi(\mathcal{U})
 \end{array} \quad (12)$$

be a local trivialization of l.a.b. (2) with $\xi : \mathcal{U} \rightarrow \xi(\mathcal{U})$ an admissible chart on M and $\bar{A}$ the standard fibre.

If we apply the functor corresponding to l.a.b. isomorphism (12) to an admissible chart $(\eta, \mathcal{W})$ on P , we obtain the diagram

$$\begin{array}{ccccc}
 & A_\xi \mathcal{W} & \xrightarrow{A_\xi \eta} & A_\xi K & \\
 \alpha_\mathcal{W}^s \swarrow & & \searrow \alpha_\mathcal{W}^t & & \searrow \alpha_K^t \\
 & \mathcal{W} & \xrightarrow{\eta} & K & \\
 \downarrow \xi & \downarrow \xi_\mathcal{W}^A & & \downarrow \xi_K^A & \\
 \mathcal{U} & \xrightarrow{\xi} & \mathcal{U} & & \\
 \downarrow \xi & \downarrow \varepsilon_\mathcal{W}^s & \downarrow \varepsilon_\mathcal{W}^t & \downarrow \varepsilon_K^s & \downarrow \varepsilon_K^t \\
 & (H \times \bar{A})\mathcal{W} & \xrightarrow{(H \times \bar{A})\eta} & (H \times \bar{A})K & \\
 \downarrow \xi & \downarrow \xi & \downarrow \xi & \downarrow \xi & \downarrow \xi \\
 H & \xrightarrow{\xi} & H & & K
 \end{array} \quad (13)$$

where, for the sake of simplicity, we have set $H = \xi(\mathcal{U})$ and $K = \eta(\mathcal{W})$.

From diagram (13) we can select the subdiagram

$$\begin{array}{ccc}
 A_\xi \mathcal{W} & \xrightarrow{A_{(\xi, \eta)}} & (\xi(\mathcal{U}) \times \bar{A})\eta(\mathcal{W}) \\
 \alpha_\mathcal{W}^s \swarrow & & \searrow \varepsilon_{\eta(\mathcal{W})}^s \\
 & \mathcal{W} & \xrightarrow{\eta} \eta(\mathcal{W}) \\
 \downarrow \xi & & \downarrow \xi \\
 \mathcal{U} & \xrightarrow{\xi} & \xi(\mathcal{U})
 \end{array} \quad (14)$$

where

$$A_{(\xi, \eta)} := \xi_{\eta(\mathcal{W})}^A \circ A_\xi \eta$$

is a bijection, which, owing to Lemma 4, is a chart of AP .

Proposition 5. *AP is a differential manifold.*

Proof. AP will be a differential manifold modelled on the affine space $(E \times \bar{A})G$ provided that the charts $A_{(\xi, \eta)}$ are C^∞ -compatible.

Without loss of generality we will consider charts ξ' and η' on M and P having the same domains $\mathcal{U}$ and $\mathcal{W}$ as ξ and η , respectively. It follows that $A_\xi = A_{\xi'}$.

We have to prove that the composition

$$\begin{aligned}
 A_{(\xi', \eta')} \circ A_{(\xi, \eta)}^{-1} &= \xi_{\eta'(\mathcal{W})}^A \circ A_{\xi' \eta'} \circ (A_\xi \eta)^{-1} \circ (\xi_{\eta(\mathcal{W})}^A)^{-1} \\
 &= \xi_{\eta'(\mathcal{W})}^A \circ A_\xi (\eta' \circ \eta^{-1}) \circ (\xi_{\eta(\mathcal{W})}^A)^{-1} : (\xi(\mathcal{U}) \times \bar{A})\eta(\mathcal{W}) \longrightarrow (\xi'(\mathcal{U}) \times \bar{A})\eta'(\mathcal{W})
 \end{aligned} \quad (15)$$

is differentiable.

(i) Let us consider the transition function obtained from (12) and the analogous local trivialization of (2) corresponding to ξ' :

$$\begin{array}{ccc} \xi(\mathcal{U}) \times \bar{A} & \xrightarrow{\xi'^A \circ (\xi^A)^{-1}} & \xi'(\mathcal{U}) \times \bar{A} \\ pr_1 \downarrow & & \downarrow pr_1 \\ \xi(\mathcal{U}) & \xrightarrow{\xi' \circ \xi^{-1}} & \xi'(\mathcal{U}) \end{array} \quad (16)$$

It is a differentiable l.a.b. morphism. It follows that for all $\xi(x) \in \xi(\mathcal{U})$ the induced mapping

$$\xi'^A|_{A_x} \circ (\xi^A|_{A_x})^{-1} : \xi(x) \times \bar{A} \longrightarrow \xi'(x) \times \bar{A} : (\xi(x), a) \mapsto (\xi' \circ \xi^{-1}(\xi(x)), \Xi^{\xi(x)}(a))$$

determines a local algebra automorphism

$$\Xi^{\xi(x)} : \bar{A} \longrightarrow \bar{A} \quad (17)$$

and, moreover, that the mapping $\xi(x) \mapsto \Xi^{\xi(x)}$ is differentiable.

Owing to Proposition 7 in [2], when we apply the functor corresponding to (17) to $\eta'(\mathcal{W})$ we get the following map:

$$\Xi_{\eta'(\mathcal{W})}^{\xi(x)} : \bar{A}\eta'(\mathcal{W}) \longrightarrow \bar{A}\eta'(\mathcal{W}) : \bar{u} \mapsto \Xi^{\xi(x)} \circ \bar{u}. \quad (18)$$

The above mapping is differentiable since, by Proposition 9 in [2], for any choice of a Cartesian coordinate system $(y^1, \dots, y^m)$ in G , it admits the following global coordinate expression

$$\begin{aligned} \eta'(\mathcal{W}) \times \underbrace{(I_{\bar{A}} \times \dots \times I_{\bar{A}})}_{m \text{ times}} &\longrightarrow \eta'(\mathcal{W}) \times \underbrace{(I_{\bar{A}} \times \dots \times I_{\bar{A}})}_{m \text{ times}} \\ (\eta'(p), (\bar{u}^j)_{j=1, \dots, m}) &\mapsto (\eta'(p), (\Xi^{\xi(x)}(\bar{u}^j))_{j=1, \dots, m}), \end{aligned} \quad (19)$$

where we have set $\bar{u}^j = \bar{u}(y^j) - \bar{u}(\eta'(p))$ for all $j = 1, \dots, m$.

(ii) Now we can prove that the mapping (15) is differentiable. It acts as follows:

$$\begin{array}{ccc} v & \xrightarrow{(\xi_{\eta(\mathcal{W})}^A)^{-1}} & (\xi^A|_{A_x})^{-1} \circ v \\ & \xrightarrow{A_{\xi}(\eta' \circ \eta^{-1})} & (\xi^A|_{A_x})^{-1} \circ v \circ (\eta' \circ \eta^{-1})^* \\ & \xrightarrow{\xi'^A_{\eta'(\mathcal{W})}} & \xi'^A|_{A_x} \circ (\xi^A|_{A_x})^{-1} \circ v \circ (\eta' \circ \eta^{-1})^* \end{array} \quad (20)$$

Owing to (9), we can use the following identification

$$\begin{aligned} (\xi(\mathcal{U}) \times \bar{A})\eta(\mathcal{W}) &\cong \xi(\mathcal{U}) \times \bar{A}\eta(\mathcal{W}) \\ v &\cong (\xi(x), \bar{v}) \end{aligned}$$

where $\bar{v} = pr_2 \circ v$. With the above identification, (20) becomes

$$\begin{aligned} \xi(\mathcal{U}) \times \bar{A}\eta(\mathcal{W}) &\longrightarrow \xi'(\mathcal{U}) \times \bar{A}\eta'(\mathcal{W}) \\ (\xi(x), \bar{v}) &\longrightarrow (\xi'(x), \Xi^{\xi(x)} \circ \bar{v} \circ (\eta' \circ \eta^{-1})^*) \\ &= \left(\xi' \circ \xi^{-1}(\xi(x)), \Xi_{\eta'(\mathcal{W})}^{\xi(x)}[\bar{A}(\eta' \circ \eta^{-1})(\bar{v})] \right) \end{aligned}$$

Owing to Proposition 10 in [2] and (i), we conclude that (15) is a differentiable mapping. $\blacksquare$

Up to now we have proved that AP is a differential manifold.

The next step is to show that both α_P^s and α_P^t are surjective submersions.

Proposition 6. *The diagram*

$$\begin{array}{c} AP \\ \downarrow \alpha_P^s \\ M \end{array}$$

is a surjective submersion.

Proof. From diagram (14) we select the subdiagram

$$\begin{array}{ccc} A_\xi \mathcal{W} & \xrightarrow{A_{(\xi, \eta)}} & (\xi(\mathcal{U}) \times \bar{A})\eta(\mathcal{W}) = \xi(\mathcal{U}) \times \bar{A}\eta(\mathcal{W}) \\ \alpha_\mathcal{W}^s \swarrow & & \searrow \varepsilon_{\eta(\mathcal{W})}^s = pr_1 \\ \mathcal{U} & \xrightarrow{\xi} & \xi(\mathcal{U}) \end{array}$$

It is the coordinate expression of α_P^s , which is a projection, and hence a surjective submersion. ■

Proposition 7. *The diagram*

$$\begin{array}{c} AP \\ \downarrow \alpha_P^t \\ P \end{array}$$

is a surjective submersion.

Proof. From diagram (14) we select the subdiagram

$$\begin{array}{ccc} A_\xi \mathcal{W} & \xrightarrow{A_{(\xi, \eta)}} & (\xi(\mathcal{U}) \times \bar{A})\eta(\mathcal{W}) = \xi(\mathcal{U}) \times \bar{A}\eta(\mathcal{W}) \\ \searrow \alpha_\mathcal{W}^t & & \searrow \varepsilon_{\eta(\mathcal{W})}^t = \bar{\alpha}_P^t \circ pr_2 \\ \mathcal{W} & \xrightarrow{\eta} & \eta(\mathcal{W}) \end{array}$$

It is the coordinate expression of α_P^t , which is a projection, and hence a surjective submersion. ■

Theorem 8. *Diagram*

$$\begin{array}{ccc} & AP & \\ \alpha_P^s \swarrow & & \searrow \alpha_P^t \\ & M & P \end{array}$$

is a double bundle.

Proof. We only have to prove part (ii) of Definition 1.

Let $(\mathcal{U}, \xi)$ be a chart around a point x of M and $(\mathcal{W}, \eta)$ a chart around a point p of P . We have

$$\begin{aligned}
(\alpha_P^s)^{-1}(\mathcal{U}) \cap (\alpha_P^t)^{-1}(\mathcal{W}) &= \left(\bigcup_{x \in \mathcal{U}} A_{(x, \cdot)} P \right) \cap \left(\bigcup_{p \in \mathcal{W}} A_{(\cdot, p)} P \right) \\
&= \left(\bigcup_{x \in \mathcal{U}} \bigcup_{p \in P} A_{(x, p)} P \right) \cap \left(\bigcup_{p \in \mathcal{W}} \bigcup_{x \in M} A_{(x, p)} P \right) \\
&= \bigcup_{x \in \mathcal{U}} \bigcup_{p \in \mathcal{W}} A_{(x, p)} P = \bigcup_{x \in \mathcal{U}} \bigcup_{p \in \mathcal{W}} A_{(x, p)} \mathcal{W} \\
&= \bigcup_{x \in \mathcal{U}} A_{(x, \cdot)} \mathcal{W} = (\alpha^{-1}(\mathcal{U})) \mathcal{W} \\
&= A_\xi \mathcal{W}.
\end{aligned}$$

Therefore diagram (14) is a double local trivialization of (α_P^s, α_P^t) . ■

2.1 α -jets of mappings.

In this section we present the connection between the jets of mappings of Ehresmann type and a special class of α -jets.

Let $x \in M$. In the algebra $C^\infty(M)$ consider the maximal ideal $I_0(M, x)$ of all functions vanishing at x , and its powers

$$I^k(M, x) = (I_0(M, x))^{k+1} \quad \forall k \in \mathbb{N}.$$

We can use these ideals in two ways.

(i) Consider the space $C^\infty(M|P)$ of all differentiable mappings from M to P and introduce, in the Cartesian product $C^\infty(M|P) \times M$, the following equivalence relation:

$$(\varphi', x') \sim_k (\varphi, x) \iff \begin{cases} x' = x \\ g \circ \varphi' - g \circ \varphi \in I^k(M, x) \end{cases} \quad \forall g \in C^\infty(P).$$

The equivalence class of (φ, x) is denoted by $j^k \varphi(x)$ and called the k -jet of φ at x . The set of all k -jets is denoted by $J^k(M|P)$.

(ii) Consider the special case of $P = \mathbb{R}$. For each $x \in M$, the set of k -jets of functions on M at x , denoted by $A_x^k(M)$, can be identified (cf. [7], Remark 12.7) with the local algebra

$$A_x^k(M) = \mathbb{R} + I_0(M, x) / I^k(M, x),$$

and then we have the local algebra bundle

$$\begin{array}{ccc}
A^k(M) & & \\
\downarrow \alpha^k & & \\
M & &
\end{array} \tag{21}$$

Now, if we apply the functor associated with (21) to differential manifold P , we obtain the double bundle

$$\begin{array}{ccc}
& A^k(M)P & \\
(\alpha^k)_P^s \swarrow & & \searrow (\alpha^k)_P^t \\
M & & P
\end{array}$$

Constructions (i) and (ii) are, actually, closely related, as shown in the following theorem.

Theorem 9. *The mapping*

$$\chi : J^k(M|P) \longrightarrow A^k(M)P$$

defined by

$$\chi(j^k \varphi(x)) : g \mapsto j^k(g \circ \varphi)(x)$$

is a bijection.

Proof. Note that χ is well defined by definition of $J^k(M|P)$. We prove bijectivity by showing that for every $u \in A^k(M)P$ there is a unique $j^k \varphi(x) \in J^k(M|P)$ such that

$$\chi(j^k \varphi(x)) = u.$$

Let $p = (\alpha^k)_P^t(u)$ and η be any chart on P around p . Given a Cartesian frame ϕ on the n -dimensional affine space G centred at $\eta(p)$, put

$$\zeta = \phi \circ \eta$$

and

$$\zeta^h = x^h \circ \zeta,$$

where x^h is the h -th natural coordinate function on $\mathbb{R}^n$. For all indices h let $\varphi^h \in C^\infty(M)$ be such that

$$u(\zeta^h) = j^k \varphi^h(x).$$

Then, owing to (3),

$$\varphi^h(x) = \zeta^h(p).$$

Up to replacing φ^h with another representative of the same jet, we may assume that φ^h is not constant. Since M is connected, it follows that the image of φ^h is an interval in $\mathbb{R}$. As a consequence the image of

$$\varphi_\zeta = (\varphi^1, \dots, \varphi^n)$$

contains an open neighborhood of

$$\zeta(p) = \varphi_\zeta(x).$$

The composition $\zeta^{-1} \circ \varphi_\zeta$ defines a mapping $\varphi \in C^\infty(M|P)$ such that

$$\varphi^h = \zeta^h \circ \varphi.$$

We will prove that the mapping φ is a representative of the required jet in $J^k(M|P)$.

Let $g \in C^\infty(P)$, and let $g_\zeta = g \circ \zeta^{-1}$. Owing to the local character of u , we have

$$u(g) = u(g_\zeta \circ \zeta).$$

We will write, for g_ζ , the Taylor formula at $\zeta(p) = 0$, of order k

$$g_\zeta = \mathcal{P}^k(x^1, \dots, x^n) + R^k$$

with Lagrange remainder

$$R^k = \mathcal{P}^{k+1}H,$$

where $\mathcal{P}^{k+1}$ is a homogeneous polynomial of degree $k+1$ at 0.

Then we have

$$\begin{aligned} g_\zeta \circ \zeta &= \mathcal{P}^k(x^1 \circ \zeta, \dots, x^n \circ \zeta) + R^k \circ \zeta \\ &= \mathcal{P}^k(\zeta^1, \dots, \zeta^n) + (\mathcal{P}^{k+1}H) \circ \zeta. \end{aligned}$$

As a consequence

$$u(g) = u(\mathcal{P}^k(\zeta^1, \dots, \zeta^n)) + u(\mathcal{P}^{k+1} \circ \zeta)u(H \circ \zeta),$$

where

$$u(\mathcal{P}^{k+1} \circ \zeta) = u(\mathcal{P}^{k+1}(\zeta^1, \dots, \zeta^n)) = \mathcal{P}^{k+1}(u(\zeta^1), \dots, u(\zeta^n)) = 0 \quad (22)$$

since $u(\zeta^h) \in I_0(M, x)/I^k(M, x)$. Therefore

$$\begin{aligned} u(g) &= \mathcal{P}^k(u(\zeta^1), \dots, u(\zeta^n)) \\ &= \mathcal{P}^k(j^k \varphi^1(x), \dots, j^k \varphi^n(x)) \\ &= j^k(\mathcal{P}^k(\varphi^1, \dots, \varphi^n))(x) \\ &= j^k(\mathcal{P}^k(\zeta^1 \circ \varphi, \dots, \zeta^n \circ \varphi))(x) \\ &= j^k(\mathcal{P}^k(\zeta^1, \dots, \zeta^n) \circ \varphi)(x) \\ &= j^k(\mathcal{P}^k(\zeta^1, \dots, \zeta^n) \circ \varphi)(x). \end{aligned}$$

Due to the local character of jets, we have

$$\begin{aligned} u(g) &= j^k(\mathcal{P}^k(\zeta^1, \dots, \zeta^n) \circ \varphi)(x) \\ &= j^k(g_\zeta \circ \zeta \circ \varphi - R^k \circ \zeta \circ \varphi)(x) \\ &= j^k(g \circ \varphi)(x) - j^k(R^k \circ \zeta \circ \varphi)(x) \\ &= j^k(g \circ \varphi)(x). \end{aligned}$$

The last equality is justified by arguments similar to those used in (22).

The uniqueness of $j^k \varphi(x)$ trivially follows by definition. ■

References

- [1] M. Barile, F. Barone and W.M. Tulczyjew, *Polarizations and Differential Calculus in Affine Spaces*, Linear Multilinear Algebra. **55**, 121–146 (2007).
- [2] M. Barile, F. Barone and W.M. Tulczyjew, *The category of local algebras and points proches*, Bull. Math. Soc. Sci. Math. R. S. Roumanie (N.S.), **50**, 3–31 (2007).
- [3] E. Dubuc, *Sur les modèles de la géométrie différentielle synthétique*, Cah. Topol. Géom. Différ. Catég., **XX**, 234–278 (1979).
- [4] E. Dubuc and J. Zilber, *Weil Prolongations of Banach Manifolds in an Analytic Model of SDG*, Cah. Topol. Géom. Différ. Catég., **XLVI**, 83–98 (2005).

- [5] D.J. Eck, *Product-Preserving Functors on Smooth Manifolds*, J. Pure Appl. Algebra, **42**, 133–140 (1986).
- [6] C. Ehresmann, *Les prolongements d'une variété différentiable*, C.R. Acad. Sci., **233**, 598–600, 777–779 (1951).
- [7] I. Kolář, P.W. Michor and J. Slovák, *Natural Operations in Differential Geometry*, Springer (1993).
- [8] I. Kolář, *Bundle functors of the jet type*, in: I. Kolář et al. (eds.), Differential geometry and applications. Proceedings of the 7th international conference, DGA 98, and satellite conference of ICM in Berlin, Brno, Czech Republic, August 10-14, 1998. Brno: Masaryk University, 231–237 (1999).
- [9] P. Lieberman, *Introduction to the theory of semi-holonomic jets*, Arch. Math. Brno, **33**, 173–189 (1996).
- [10] G. Pidello, W.M. Tulczyjew, *Derivations of differential forms on jet bundles*, Ann. Mat. Pura Appl., IV. Ser., **CXLVII**, 249–352 (1987).
- [11] J. Pradines, *Fibrés vectoriels doubles et calcul des jets non holonomes*, Esquisses Math., **29**, Amiens (1977).
- [12] V.V. Shurygin, *The structure of smooth mappings over Weil algebras and the category of manifolds over algebras*, Lobachevskii J. Math., **5**, 29-55 (1999).
- [13] W.M. Tulczyjew, *Evolution of Ehresmann's jet theory*. To appear in: Banach Center Publications, Institute of Mathematics, Polish Academy of Sciences, Warszawa.
- [14] A. Weil, *Théorie des points proches sur les variétés différentiables*, Colloques internat. Centre nat. Rech. Sci., **52**, 111–117 (1953).